

# New physics effects on the $hZZ$ and $hhh$ vertices

Shinya Kanemura  
(Osaka Univ)

Y.Okada, E.Senaha (KEK), C.-P.Yuan (Michigan SU)

3. October 2003 at 2nd LC Physics Study Group Meeting

- I. Introduction
- II. The one-loop  $hZZ$  and  $hhh$  vertices in the two Higgs doublet model
- III. Numerical results
- IV. Summary

# I. Introduction

The Higgs boson  $\implies$  yet to be confirmed

$m_h$ : free in the SM

- Theoretical bounds:

$$140 \lesssim m_h^{\text{SM}} \lesssim 175 \text{ GeV for } \Lambda = 10^{19} \text{ GeV}$$

(In MSSM,  $m_h < 120\text{-}130 \text{ GeV}$ )

- Experimental bounds:

LEP data:  $114 \text{ GeV} < m_h < 196 \text{ GeV}$  (SM)

$h$  will be discovered at **Tevatron, LHC**



Once a Higgs boson ( $h$ ) is found,  
a **precision study** of Higgs couplings

$$hVV, h\bar{f}f, hhh$$

is done at a **Linear Collider (LC)**

# Higgs coupling measurements at LC

- Higgs-gauge coupling  $g_{hWW}, g_{hZZ}$

⇒ Higgs Mechanism

$$\sigma(e^+e^- \rightarrow Zh), \sigma(e^+e^- \rightarrow \bar{\nu}\nu h) \Rightarrow \mathcal{O}(1\%)$$


- Yukawa coupling  $Y_f$

⇒ Mass Generation of Matter

$$B(h \rightarrow \bar{f}f') \Rightarrow \mathcal{O}(1\%)$$

- Higgs self-coupling  $\lambda_{hhh}$

⇒ Dynamics of Higgs sector



$\lambda_{hhh}$  can be measured by  $\mathcal{O}(10 - 20\%)$  accuracy

$$\text{LC: } \sqrt{s} = 0.5 - 1.5 \text{ TeV, } \mathcal{L} = 1 \text{ ab}^{-1}$$

Battaglia et al, ACFA Higgs WG

The potential for precision measurement at a LC motivates us to study radiative corrections.

$$\Delta^{\text{Exp}} g_{hVV}/g_{hVV} = \mathcal{O}(1\%)$$
$$\Delta^{\text{Exp}} \lambda_{hhh}/\lambda_{hhh} = \mathcal{O}(10 - 20\%)$$

## Leading top-loop contribution in the SM

- $hVV$  coupling

$$M_{hVV}^{\mu\nu} = M_1 g^{\mu\nu} + M_2 p_1^\nu p_2^\mu + M_3 i\epsilon^{\mu\nu\rho\sigma} p_{1\rho} p_{2\sigma}$$

$$g_{hZZ} \equiv M_1 = \frac{2m_Z^2}{v} \left( 1 - \frac{5N_c m_t^2}{96\pi^2 v^2} + \dots \right),$$

loop effects  $\sim 1\%$

- $hhh$  coupling

$$\lambda_{hhh} \sim \frac{3m_h^2}{v} \left( 1 - \frac{N_c m_t^4}{3\pi^2 v^2 m_h^2} + \dots \right)$$

loop effects  $\sim 10\%$

A  $m_t^4$  term appears in the renormalized  $\lambda_{hhh}$ .

The corrections are comparable to the measurement accuracy  
 $\Rightarrow$  **How about new physics effects?**

# Probe of new physics via Higgs sector

## New Physics

- Naturalness

SUSY, Dynamical EW breaking, Little Higgs, Extra D .....

- EW baryogenesis

- Neutrino mass

- Top-bottom mass hierarchy



## Low-energy Effective Theory

### Extended Higgs sectors

with additional doublets, singlets, ...



**Extra Higgs scalars**  $H^\pm, A, \dots$

⇒ different prediction to  $hVV, h\bar{f}f$  and  $hhh$ .



**We can determine direction of new physics  
by a detailed study of the Higgs masses and couplings**

## Property of extra scalars ( $H, H^\pm, A, \dots$ ) in extended Higgs sectors

### SM like Higgs boson ( $h$ ) mass

$$m_h^2 \sim \lambda v^2, \text{ VEV: } v (\simeq 246 \text{ GeV})$$

### Extra Higgs boson masses

$$\left. \begin{array}{l} m_H \\ m_{H^\pm} \\ m_A \\ \dots \end{array} \right\} \sim \lambda_i v^2 + M^2$$

## In this talk:

We evaluate  
the one-loop corrected  $hZZ$  and  $hhh$  vertices  
in the two-Higgs doublet model (THDM).

### THDM

- Simplest extension of the SM Higgs sector
- Typical aspects of extended Higgs sectors motivated by various new physics scenarios

How much can the deviation from  
the SM prediction be substantial  
under theoretical and experimental constraints?

# The two Higgs doublet model (THDM)

- THDM with a softly-broken discrete symmetry:

$$(\Phi_1 \rightarrow \Phi_1; \Phi_2 \rightarrow -\Phi_2)$$

$$\Rightarrow \text{Natural FCNC suppression}$$

Yukawa interaction (Model I, II):

$$\mathcal{L}_I = -y_D \bar{Q}_L \Phi_1 b_R - y_U \bar{t}_R \Phi_1^\dagger Q_L + (h.c.).$$

$$\mathcal{L}_{II} = -y_D \bar{Q}_L \Phi_1 b_R - y_U \bar{t}_R \Phi_2^\dagger Q_L + (h.c.).$$

Higgs potential:

$$V_{\text{THDM}} = +m_1^2 |\Phi_1|^2 + m_2^2 |\Phi_2|^2 - \underline{m_3^2 (\Phi_1^\dagger \Phi_2 + \Phi_2^\dagger \Phi_1)}$$

$$+ \frac{\lambda_1}{2} |\Phi_1|^4 + \frac{\lambda_2}{2} |\Phi_2|^4 + \lambda_3 |\Phi_1|^2 |\Phi_2|^2$$

$$+ \lambda_4 |\Phi_1^\dagger \Phi_2|^2 + \frac{\lambda_5}{2} \left[ (\Phi_1^\dagger \Phi_2)^2 + (h.c.) \right]$$

$\Phi_1$  and  $\Phi_2 \Rightarrow h, H, A^0, H^\pm \oplus 3$  Goldstone bosons

$\uparrow \quad \uparrow \quad \uparrow$  charged  
 CPeven CPodd

8 parameters :  $\Rightarrow \{m_h, m_H, m_A, m_{H^\pm}, \alpha, \beta, v, M_{\text{soft}}\}$

$v$  (VEV)  $\simeq 246$  GeV,  $\tan \beta (= \langle \Phi_2 \rangle / \langle \Phi_1 \rangle)$

$\alpha$ : mixing angle between  $h$  and  $H$

$M_{\text{soft}} (= \frac{m_3}{\sqrt{\cos \beta \sin \beta}})$ : soft-breaking scale  
of the discrete symm.



- Masses of physical Higgs bosons:

$$m_h^2 = v^2 \left( \lambda_1 \cos^4 \beta + \lambda_2 \sin^4 \beta + \frac{\lambda}{2} \sin^2 2\beta \right) + \mathcal{O}\left(\frac{v^2}{M_{\text{soft}}^2}\right),$$

$$m_H^2 = M_{\text{soft}}^2 + v^2 (\lambda_1 + \lambda_2 - 2\lambda) \sin^2 \beta \cos^2 \beta + \mathcal{O}\left(\frac{v^2}{M_{\text{soft}}^2}\right),$$

$$m_{H^\pm}^2 = M_{\text{soft}}^2 - \frac{\lambda_4 + \lambda_5}{2} v^2,$$

$$m_A^2 = M_{\text{soft}}^2 - \lambda_5 v^2.$$

$M_{\text{soft}}$ : soft breaking scale

of the discrete symmetry

- $M_{\text{soft}}$  determines decoupling/non-decoupling property of heavy Higgs bosons ( $\Phi = A, H^\pm$  or  $H$ )

$$m_\Phi^2 = M_{\text{soft}}^2 + \lambda_i v^2$$

Decoupling: for  $m_\Phi^2 \sim M_{\text{soft}}^2$  ( $M_{\text{soft}}^2 \gg \lambda v^2$ )

Loop-effects of  $H, A, H^\pm$  decouple

(Decoupling Theorem)

Non-Decoupling: for  $m_\Phi^2 \sim \lambda_i v^2$  ( $M_{\text{soft}}^2 \lesssim \lambda v^2$ )

$m_\Phi^n$  terms in the renormalized low energy observables

(similar to the top effects:  $m_t^2 = y_t^2 v^2$ )

# The tree-level coupling constants

## THDM

$$g_{hZZ}^{\text{tree}} = +\frac{2m_Z^2}{v} \sin(\beta - \alpha)$$

$$\lambda_{hhh}^{\text{tree}} = -\frac{3}{2v \sin 2\beta} \left[ \{ \cos(3\alpha - \beta) + 3 \cos(\beta + \alpha) \} m_h^2 - 4 \cos^2(\alpha - \beta) \cos(\alpha + \beta) M_{\text{soft}}^2 \right]$$

- $\alpha \rightarrow \beta - \pi/2$  (Decoupling Limit)

$$g_{hZZ}^{\text{tree}} \rightarrow g_{hZZ}^{\text{tree}}(\text{SM}) = \frac{2m_Z^2}{v},$$
$$\lambda_{hhh}^{\text{tree}} \rightarrow \lambda_{hhh}^{\text{tree}}(\text{SM}) = \frac{3m_h^2}{v}$$

Loop correction is essentially important!

- $\alpha \neq \beta - \pi/2$

The tree-level deviation from the SM prediction appears.

Tree-Level Deviation vs Loop Correction

## One-loop effects on $g_{hZZ}$ and $\lambda_{hhh}$

- Contributions of heavy quarks and Higgs bosons
- Renormalization: **On-shell scheme**

The leading 1-loop contribution  
in **Decoupling Limit** ( $\alpha \rightarrow \beta - \pi/2$ )


$$g_{hZZ}^{\text{loop}} = \frac{2m_Z^2}{v} \left[ 1 - \frac{1}{16\pi^2} \left\{ \frac{5N_c m_t^2}{6v^2} + \frac{2m_\Phi^2}{3v^2} \left( 1 - \frac{M_{\text{soft}}^2}{m_\Phi^2} \right)^2 \right\} \right]$$

$$\lambda_{hhh}^{\text{loop}} = -\frac{3m_h^2}{v} \left[ 1 + \frac{1}{16\pi^2} \left\{ -\frac{16N_c m_t^4}{3v^2 m_h^2} + \frac{16m_\Phi^4}{3m_h^2 v^2} \left( 1 - \frac{M_{\text{soft}}^2}{m_\Phi^2} \right)^3 \right\} \right]$$


In  $\lambda_{hhh}^{\text{loop}}$ ,  $\mathcal{O}(m_\Phi^4)$  ( $\Phi = H, A, H^+$ ) terms appear  
with a suppression factor

$$m_\Phi^4 \left( 1 - \frac{M_{\text{soft}}^2}{m_\Phi^2} \right)^3 \rightarrow \begin{cases} \frac{\lambda_i v^2}{m_\Phi^2}, & (m_\Phi^2 \sim M_{\text{soft}}^2), \\ & \text{(decoupling for } m_\Phi \rightarrow \infty) \\ m_\Phi^4, & (m_\Phi^2 \sim \lambda_i v^2), \\ & \text{(non-decoupling effect)} \end{cases}$$

# Non-decoupling Effects



*hZZ* coupling  
 Deviation  
 from the SM  
 $\sim 1\%$



*hhh* coupling  
 Deviation  
 from the SM  
 $\sim 30-100\%$   
 due to  
 the  $m_A^4$  term

$$\delta \equiv 1 - \sin^2(\alpha - \beta)$$

$$M_{\text{soft}} = 0, \tan \beta = 1, m_A = m_H = m_{H^\pm}$$

# Scan Analysis

Free Parameters in the THDM:

$$\tan \beta, m_H, m_{H^\pm}, m_A, M_{\text{soft}}.$$

for fixed  $m_h$ , and  $\delta \equiv 1 - \sin^2(\alpha - \beta)$ .

Searching allowed region of the deviation from the SM  
in the  $hZZ$  and  $hhh$  vertices,  
under the constraint from

- **Perturbative unitarity**

$$|a^0(W_L^+ W_L^- \rightarrow W_L^+ W_L^-)| < \xi, \quad (\xi = 1/4)$$

for 15 channels  $W_L^+ W_L^-, Z_L Z_L, Z_L h, hh, hH, \dots$


- **Vacuum Stability**

$$V_{\text{eff}}(\langle \Phi_1 \rangle, \langle \Phi_2 \rangle) \geq 0 \text{ for } \langle \Phi_i \rangle \rightarrow \infty.$$


- **LEP Precision Data:**

Constraint on the (S,T,U) parameters

# Allowed Region ( $M_{\text{soft}} = 0$ )





*hZZ* coupling  
Deviation  
from the SM  
 $\sim 1\%$



*hhh* coupling  
Deviation  
from the SM  
 $\lesssim 50\text{-}100\%$   
due to  
the  $m_A^4$  term

$$\delta \equiv 1 - \sin^2(\alpha - \beta)$$

# Allowed Region ( $M_{\text{soft}} = m_A/\sqrt{2}$ )



$$\delta \equiv 1 - \sin^2(\alpha - \beta)$$

# Summary

One-loop effective couplings of  $hZZ$  and  $hhh$  in the SM and the THDM

Scan Analysis:

- $hZZ$ : deviation from the SM

One-loop  $\mathcal{O}(m_A^2)$  contribution ( $\sim 1\%$ )

⊕ Tree-level difference when  $\delta = 1 - \sin^2(\alpha - \beta) \neq 0$

- $hhh$ : deviation from the SM

One-loop  $\mathcal{O}(m_A^4)$  contribution  $\lesssim 50-100\%$

⊕ Tree-level difference when  $\delta \neq 0$

Even when the  $hZZ$  measurement is consistent with the SM by  $\mathcal{O}(1)\%$ ,  $hhh$  can deviate from the SM by  $\sim 30-100\%$  due to the non-decoupling effects of heavy Higgs bosons

Such deviation can be tested at a Linear Collider