

# Luminosity Monitor and Active Mask

## (1) Luminosity monitor

geometrical acceptance:

$$163 < z < 178 \text{ cm} \quad \text{and} \quad 0.05 < \theta < 0.15 \text{ rad.}$$

made of tungsten only in this study

segmentation:

r	32 divisions	$\Delta r \sim 5\text{mm}$
$\phi$	16 divisions	$\Delta\phi \sim 3.2 - 9.7 \text{ cm}$
z	128 divisions	$\Delta z \sim 1.17\text{mm}$

## (2) Active mask (front part of conical mask)

geometrical acceptance:


$$30 < z < 37.5 \text{ cm} \quad \text{and} \quad 0.15 < \theta < 0.20 \text{ rad.}$$

made of tungsten(W) and silicon pad( Si,  $200\mu\text{m}^t$ )

$5\text{mm}^t\text{W/Si}/(1\text{cm}^t\text{W/Si})^7$  8 layers

segmentation:

r	8-10 divisions	$\Delta r = 2\text{mm}$
$\phi$	32 divisions	$\Delta\phi \sim 0.9 - 1.2 \text{ cm}$



# Generated events

(1)  $e^+e^-$  pairs by cain21d  
parameter A at  $E_{cm}=500\text{GeV}$   
100 bunch crossings

correspond to 1 train crossing

(2) an electron/ muon into luminosity monitor  
and active mask

50 GeV and 250 GeV electrons

250 GeV muons


## Simulation results

(1) Energy deposits in total volume





B=2T	$e^+e^-$ pairs	50GeV e	250GeV e	250GeV $\mu$
am	264MeV	120.4MeV 121.7MeV	541.0MeV 548.0MeV for 10 layers	0.48MeV
lm	152GeV	49.9GeV	249.5GeV	1.67GeV

B=3T	$e^+e^-$ pairs
am	29.3MeV
lm	46.7GeV


am=active mask, lm=luminosity monitor




a 50 GeV electron with 100 bunch crossings: Luminosity monitor




a 50 GeV electron with 100 bunch crossings: Luminosity monitor




+ LM:energy deposit(keV):Z vs PHI



PROY + LM:energy deposit(keV):Z vs PHI




PROX + LM:energy deposit(keV):Z vs PHI




+ LM:energy deposit(keV):Z vs PHI

a 250 GeV electron with 100 bunch crossings : Luminosity monitor




+ LM:energy deposit(keV):Z vs R




ID=231, N=325

PROY + LM:energy deposit(keV):Z vs R




ID=232, N=325

PROX + LM:energy deposit(keV):Z vs R




+ LM:energy deposit(keV):Z vs R

a 250 GeV electron with 100 bunch crossings : Luminosity monitor




+ LM:energy deposit(keV):Z vs PHI




ID=235, N=325

PROY + LM:energy deposit(keV):Z vs PHI




ID=236, N=325

PROX + LM:energy deposit(keV):Z vs PHI




+ LM:energy deposit(keV):Z vs PHI







*a 250GeV muon into active mask*




*AM:energy deposit(keV):Layer vs R*



*AM:energy deposit(keV):Layer vs phi*




*ID=490, N=8  
AM:energy deposit (KeV) as a function of Z*




*ID=491, N=8  
AM:energy flow (GeV) as a function of Z*


*a 50 GeV electron into active mask*




*AM:energy deposit(keV):Layer vs phi*



*AM:energy deposit(keV):Layer vs phi*




*+ AM:energy deposit(keV):Layer vs phi*




*+ AM:energy deposit(keV):Layer vs phi*


*a 50 GeV electron into active mask*




*ID=490,N=1777*  
*AM:energy deposit (KeV) as a function of Z*



*ID=491,N=1777*  
*AM:energy flow (GeV) as a function of Z*




*ID=1490,N=115*  
*AM:energy deposit (KeV) as a function of Z*




*ID=1491,N=115*  
*AM:energy flow (GeV) as a function of Z*


*a 250 GeV electron into active mask*




*AM:energy deposit(keV):Layer vs phi*



*AM:energy deposit(keV):Layer vs phi*




*+ AM:energy deposit(keV):Layer vs phi*




*+ AM:energy deposit(keV):Layer vs phi*


*a 250 GeV electron into active mask*




*ID=490,N=1777*  
*x 10<sup>4</sup> AM:energy deposit (KeV) as a function of Z*



*ID=491,N=1777*  
*AM:energy flow (GeV) as a function of Z*





*ID=2490,N=212*  
*AM:energy deposit (KeV) as a function of Z*




*ID=2491,N=212*  
*AM:energy flow (GeV) as a function of Z*


*TOP: B=3 tesla and BOTTOM: B=2 tesla*



*TOP: B=3 tesla and BOTTOM: B=2 tesla*



*AM: energy deposit(keV): Layer vs phi*



*AM: energy deposit(keV): Layer vs phi*

# Conclusions

## (1) Luminosity monitor

Among the total energy deposit of 152 (46.7) GeV/train due to  $e^+e^-$  pairs, only  $54^*(14^*)$  GeV comes from the front at  $B=2(3)T$ , while most comes from the inner-back.

(\* sum of incoming energies)

$\phi$  segmentation (16 div.) is very important,  $r$  segmentation is desired to determine  $\theta$  with  $\delta\theta \sim$  a few m radian.

So, a fine-segmented W/Silicon calorimeter seems to be ideal. Thickness of tungsten must be optimized in terms of energy-resolution.

## (2) Active Mask

First layer (5mm<sup>t</sup> W) has  $\sim 50\%$  energy deposit for  $e^+e^-$  pairs.

$\phi$  segmentation (32 div.) is very important.

8 layers of W/Si-pad calorimeter works very well for vetoing high energy electrons.